



Soils Map



State: **Minnesota**
 County: **Kandiyohi**
 Location: **16-119N-34W**
 Township: **Kandiyohi**
 Acres: **257.54**
 Date: **7/6/2021**



Soils data provided by USDA and NRCS.

Area Symbol: MN067, Soil Area Version: 19						
Code	Soil Description	Acres	Percent of field	Non-Irr Class Legend	Non-Irr Class *c	Productivity Index
L223B	Amiret-Swanlake loams, 2 to 6 percent slopes	67.86	26.3%		Ile	92
954C	Swanlake-Ves loams, 6 to 12 percent slopes	52.08	20.2%		IIIe	78
L186A	Danielson-Danielson, overwash complex, 1 to 4 percent slopes	31.94	12.4%		IIw	93
981	Canisteo-Harps loams	27.08	10.5%		IIw	92
L83A	Webster clay loam, 0 to 2 percent slopes	21.86	8.5%		IIw	93
446B	Normania loam, 2 to 5 percent slopes	19.31	7.5%		Ile	94
899	Harps-Okoboji complex, 0 to 2 percent slopes	7.34	2.9%		IIw	89
L166C2	Newlondon-Strout complex, 6 to 12 percent slopes, moderately eroded	6.97	2.7%		IIIe	77
954D	Swanlake-Ves loams, 12 to 18 percent slopes	6.52	2.5%		IVe	61
110	Marna silty clay loam, 0 to 2 percent slopes	5.90	2.3%		IIw	87
L163A	Okoboji silty clay loam, 0 to 1 percent slopes	3.57	1.4%		IIIw	86
L33B	Kandiyohi clay, 2 to 5 percent slopes	2.83	1.1%		Ile	90
L185B	Strout-Arkton complex, 2 to 6 percent slopes	2.01	0.8%		Ile	83
L164A	Lura silty clay, depressionnal, firm substratum, 0 to 1 percent slopes	1.45	0.6%		IIIw	81
817	Canisteo-Seaforth complex, 0 to 3 percent slopes	0.82	0.3%		IIw	94
Weighted Average						87.9

*c: Using Capabilities Class Dominant Condition Aggregation Method
 Soils data provided by USDA and NRCS.